



## King's Research Portal

DOI:

[10.1038/s41366-020-00682-5](https://doi.org/10.1038/s41366-020-00682-5)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Maragkoudaki, X., Naylor, M., Papacleovoulou, G., Stolarczyk, E., Rees, D., Pombo, J. M., Abu-Hayyeh, S., Czajka, A., Howard, J. K., Malik, A. N., Williamson, C., Poston, L., & Taylor, P. D. (2020). Supplementation with a prebiotic (polydextrose) in obese mouse pregnancy improves maternal glucose homeostasis and protects against offspring obesity. *International Journal of Obesity*, 44(12), 2382-2393. <https://doi.org/10.1038/s41366-020-00682-5>

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### Take down policy

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

SUPPLEMENTATION WITH A PREBIOTIC (POLYDEXTROSE) IN  
OBESE MOUSE PREGNANCY IMPROVES MATERNAL GLUCOSE  
HOMEOSTASIS AND PROTECTS AGAINST OFFSPRING OBESITY

**Authors:** Xanthi Maragkoudaki<sup>1</sup>, Matthew Naylor<sup>1</sup>, Georgia Papacleovoulou<sup>1</sup>, Emilie Stolarczyk<sup>2</sup>, Douglas Rees<sup>1</sup>, Joaquim M. Pombo<sup>1</sup>, Shadi Abu-Hayyeh<sup>1</sup>, Elena Zambrano<sup>2</sup>, Anja Czajka<sup>3</sup>, Jane K. Howard<sup>3</sup>, Afshan N. Malik<sup>3</sup>, Catherine Williamson<sup>1</sup>, Lucilla Poston<sup>1</sup> & Paul D. Taylor<sup>1</sup>

**Institution:** <sup>1</sup>Department of Women and Children's Health, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London. <sup>2</sup>Reproductive Biology Department, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, México. <sup>3</sup>Department of Diabetes Research, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London.

**Corresponding Author:**

Dr Paul Taylor  
Department of Women and Children's Health  
School of Life Course Sciences,  
Faculty of Life Sciences and Medicine  
King's College London  
10th Floor, North Wing.  
St Thomas' Hospital  
1 Westminster Bridge  
SE1 7EH  
Tel. +44 (0) 20 7188 3630  
Fax. +44 (0) 20 7620 1227  
Email. Paul.taylor@kcl.ac.uk

**Subject Terms:** Obesity, pregnancy, mouse. appetite, energy expenditure, prebiotic, polydextrose, developmental programming.

**Conflict of Interest Statement:** The authors declare no conflict of interest.

**Word Count 3964/4000**

### ABSTRACT

**Objectives:** We hypothesised that maternal diet-induced-obesity has adverse consequences for offspring energy expenditure and susceptibility to obesity in adulthood, and that the prebiotic polydextrose (PDX) will prevent the consequences of programming by maternal obesity.

**Methods:** Female mice were fed a control (Con), or obesogenic diet (Ob) for 6 weeks prior to mating and throughout pregnancy and lactation. Half the obese dams were supplemented with 5% PDX (ObPDX) in drinking water throughout pregnancy and lactation. Offspring were weaned onto standard chow. At 3 and 6 months, offspring energy intake (EI) and energy expenditure (EE by indirect calorimetry) were measured, and a glucose-tolerance-test performed. Offspring of control (OffCon), obese (OffOb) and PDX supplemented (OffObP) dams were subsequently challenged for 3-weeks with Ob, and energy balance reassessed. Potential modifiers of offspring energy balance including gut microbiota and biomarkers of mitochondrial activity were also evaluated.

**Results:** 6-month-old male OffOb demonstrated increased body weight (BW,  $P<0.001$ ) and white adipose tissue mass ( $P<0.05$ ), decreased brown adipose tissue mass (BAT,  $P<0.01$ ), lower night-time EE ( $P<0.001$ ) *versus* OffCon, which were prevented in OffObP. Both male and female OffOb showed abnormal GTT (peak [Glucose]  $P<0.001$ ; AUC,  $P<0.05$ ) which was prevented by PDX. The Ob challenge resulted in greater BW gain in both male and female OffOb *versus* OffCon ( $P<0.05$ ), also associated with increased EI ( $P<0.05$ ) and reduced EE in females ( $P<0.01$ ). OffObP were protected from accelerated BW gain on the OB diet compared with controls, associated with increased night-time EE in both male ( $P<0.05$ ) and female OffObP ( $P<0.001$ ). PDX also prevented an increase in skeletal muscle mtDNA copy number in OffOb *vs.* OffCon ( $P<0.01$ ) and increased the percentage of Bacteroides cells in faecal samples from male OffObP relative to controls.

**Conclusions:** Maternal obesity adversely influences adult offspring energy balance and propensity for obesity, which is ameliorated by maternal PDX-treatment with associated changes in gut microbiota composition and skeletal muscle mitochondrial function.

## INTRODUCTION

Maternal obesity constitutes the most common obstetric risk factor in developed countries with direct implications not only for maternal and neonatal morbidity and mortality but also for increased risk of obesity in the next generation (1-3). Mother-child cohort studies suggest the acquisition of obesogenic traits from mother via an undefined association between maternal body mass index (BMI) in pregnancy and risk of obesity in childhood and beyond (2). Increasing experimental evidence suggests that exposure to maternal obesity *in utero* and during lactation, especially maternal hyperglycaemia and insulin resistance (4) associated inflammation and metabolic dysfunction, may contribute to this relationship (5), impacting Global Sustainable Development Goals, in terms of health and wellbeing of current and future generations (6). Interventions are therefore urgently sought.

In view of this unmet clinical need, we have investigated the potential of a dietary supplement to improve the maternal metabolic profile in obese pregnant mice and thereby prevent deleterious effects on offspring metabolism, inflammation and energy balance. Polydextrose (PDX) is a low calorie, neutral tasting, condensation polymer of D-glucose, sorbitol, and citric acid, which is water soluble, resistant to digestion in the small intestine, but partially fermented by endogenous microbiota in the large intestine, leading to its classification as a soluble dietary fibre (7). Randomised placebo-controlled trials and two recent meta-analysis of studies in adult humans have reported increased satiety,

and improved glucose homeostasis and lipid profiles with PDX supplementation (7-11). Therefore, PDX supplementation in obese women offers the potential to improve metabolic profile and inflammation during pregnancy to positively impact on the developing offspring (12, 13).

We have previously reported cardiometabolic dysfunction in the offspring of mice with diet-induced obesity (14-17). In this study we have addressed the effect of obesity and PDX supplementation on offspring metabolic function, with a focus on energy balance, both intake and expenditure. Energy expenditure has been relatively under-explored, in models of maternal/offspring obesity. A recent meta-analysis addressing the effect of maternal obesogenic diets in rodents on offspring food intake and body mass concluded that, overall, effects on appetite are modest, whereas the increase in offspring body weight are consistent with permanent alterations in metabolism (18).

## **MATERIALS AND METHODS**

### ***Animal husbandry***

All studies were approved locally by the Animal Welfare and Ethics Committee (AWERB) and were conducted under UK Home Office License (Taylor, PPL 70/7090). Power calculations were performed based on previous in vivo data to estimate sample size. Female C57BL/6J mice were fed either a standard chow diet (RM1, Special Dietary Services, UK) or a semisynthetic obesogenic diet (approx. 16% fat, 33% simple sugars, 15% protein, total energy 16.7 kJ/g (4.0 kcal/g), as previously described (14) (Supplementary methods and Supplementary Table 1). Following successful mating, a sub-group of obesogenic diet-fed dams were randomly assigned to supplementation with PDX (5% w/v) in the drinking water throughout gestation and lactation (n=34), generating three experimental groups; control (Con); obese (Ob) and obese + PDX, (ObP,

Figure 1). This concentration of PDX has previously proven efficacious in reducing insulin resistance in adult non-pregnant rats, without adverse effects or alteration in calorific intake (19).

Offspring of Control dams (OffCon), Obese dams (OffOb) and Obese PDX supplemented dams (OffObP) were weaned and maintained on standard chow, and one male and one female from each litter studied at time points 30 days, 3 and 6 months of age. Therefore, no evaluation included more than one subject of each sex from each litter

### ***Indirect Calorimetry***

Energy expenditure (EE), respiratory exchange ratio (RER) and food intake in the offspring, were measured using LabMaster<sup>®</sup> Automated Home Cage Phenotyping (TSE Systems, Bad Homburg, Germany).

### ***Organ Collection***

At each time point, animals were killed by rising concentration of CO<sub>2</sub> or cervical dislocation, in accordance with Schedule 1 of UK Home Office guidelines. All animals were sacrificed mid-morning, blood was taken by cardiac puncture, organs were removed and immediately snap frozen in liquid nitrogen for deoxyribonucleic acid (DNA) extraction and the fat pads (perineal, gonadal, inguinal and subcutaneous) and the skeletal muscle *tibialis anterior* were weighed.

### ***Glucose tolerance test***

PDX has been shown to improve glucose tolerance in mice (20). To determine whole body glucose tolerance, an intra-peritoneal glucose tolerance test (ipGTT) was performed in the dams at gestational day 16 (GD16) and in the offspring at 30, 90 and 180 days of age. Animals were injected (i.p.) with a glucose load (1 g/kg; 10% glucose solution). Blood

glucose was measured at 15, 30, 60 and 120 minutes after glucose injection using an AlphaTRAK® Glucose meter (Abbott Animal Health).

### **Cytokine Profile**

To assess the impact of PDX on inflammatory cytokines in obese pregnancy, a subgroup of dams (n=5) were killed at gestational day 16, by a rising concentration of CO<sub>2</sub> and maternal blood samples were taken by cardiac puncture and serum stored at -80 °C. Twenty-four adipocytokines were measured from pooled serum samples, using a Proteome Profiler Mouse Adipokine Array kit (R&D Systems) (see Supplementary methods for details).

### ***Obesogenic dietary challenge***

In a separate cohort of offspring (OffCon, OffOb and OffObP) at 3 months of age, 1 male and 1 female from each litter were provided *ad libitum* access to the maternal obesogenic diet (see above) for three weeks, to assess the impact of an obesogenic dietary challenge on the adult phenotype.

### ***Quantitative real-time PCR***

Greater brown fat distribution and activation may influence energy expenditure due to increased metabolic activity. We therefore evaluated expression of relevant brown fat genes (see Supplementary Table S2 for primers and sequences). Total RNA was extracted from BAT samples with the RNeasy mini kit (QIAGEN). RNA (1 µg) was reverse transcribed into cDNA with the Superscript II kit (Invitrogen). Semi-quantitative real-time PCR with SYBR Green JumpStart Taq ReadyMix (Sigma-Aldrich) was used to detect and amplify target cDNA. Relative gene expression was calculated using the  $\Delta\Delta$  Ct method. Genes of interest were normalised to the housekeeping gene Cyclophilin B.

177

178 **Mitochondrial DNA Copy Number in Offspring Skeletal Muscle**

179 MtDNA content varies between different cell types depending on the bioenergetic  
 180 needs, but can also change in response to physiological stimuli, leading to  
 181 alterations of mtDNA being employed as a biomarker of mitochondrial  
 182 dysfunction (21, 22). Total genomic DNA was isolated from skeletal muscle using the  
 183 DNeasy blood and tissue kit (Qiagen, UK) according to the manufacturer's  
 184 guidelines, and treated by sonication to minimise effects of dilution bias. Absolute  
 185 mtDNA copy number was determined by real time qPCR. The primers (see  
 186 Supplementary Table 2 for sequences) used were specific to mouse  
 187 mitochondrial and nuclear genome targets (mMitoF1/R1 and mB2MF1/R1  
 188 respectively), as detailed previously (22).

189

190 **Analysis of offspring faecal microbiota**

191 Since PDX is hypothesised to influence the maternal microbiome (13) with vertical  
 192 transfer to neonates, we investigated broad spectrum faecal microbiota profiles in  
 193 offspring at weaning, 3 months and 6 months of age. Offspring faecal samples were snap  
 194 frozen and stored at -80°C. Samples were quantified for broad-spectrum gut bacterial  
 195 species using probes targeting six phylogenetic groups (for detailed methods, targets and  
 196 specific probes see Supplementary Methods). Phylogenetic characterisation was  
 197 performed using 16S rRNA *in situ* hybridisation and whole cell fluorescence *in situ*  
 198 hybridisation (FISH) combined with flow cytometry as described by Rigottier-Gois and  
 199 colleagues (23) employing 16S rRNA-targeted oligonucleotide probes, and targets for  
 200 rRNA dot-blot hybridisation (Panel of group- and species-specific 16S rRNA-targeted  
 201 oligonucleotide probes, Supplementary Table S4).



### **Statistical Analysis**

Data are expressed as means  $\pm$  SEM. Statistical analysis was performed with GraphPad Prism 5, (GraphPad Software Inc. San Diego, California, USA). When comparing more than two groups, one-way ANOVA followed by Bonferroni post hoc test was employed. When comparing two groups, Student's t-tests was used. Normal distributions and equality of variance between groups were checked by visual inspection of scatter plots. Statistical significance was considered when P value  $<0.05$ .  $\chi^2$  test was used to test differences in reproductive outcomes between experimental groups.

## **RESULTS**

### **Maternal Characteristics**

#### ***Body weight, food intake in Pregnancy***

There was no difference in gestational weight gain or calorific intake during gestation between the obese dams and the obese dams supplemented with PDX (Figure 2A, 2B).

#### ***Glucose Tolerance in Pregnancy***

The obese dams receiving PDX demonstrated improved glucose tolerance (Figure 2C) and a reduced area under the glucose curve (AUC) 2 hours after the i.p. glucose load compared to the obese dams GD16 (Figure 2C).

#### ***Reproductive success***

Maternal obesity affected both fertility and pup survival rates and was associated with increased rates of cannibalism in the obese dams. Control dams had 89% successful pregnancies and only 6% cannibalization compared to 44 % and 18 % respectively, for obese dams. Administration of PDX in obese pregnant and lactating dams improved

fertility rates by 14% and reduced cannibalization of the newborn pups ( $P < 0.05$ , Chi-squared test, data not shown).

### ***Maternal Cytokine profile at Gestational day 16***

Inflammatory markers were decreased in obese dams following PDX dietary supplementation; notably, TNF- $\alpha$  and CSF-1 showed a 4 and 3 -fold decrease respectively (figure 2F).

### **Offspring Characteristics**

#### ***Birth weight and litter size***

There was no influence of maternal obesity or PDX on the birth weight of offspring. There was a reduction in the litter size due to maternal obesity, which was partially reversed by maternal dietary supplementation with PDX during pregnancy (figure 2D and 2E).

#### ***Body Composition, Energy Balance and Glucose Tolerance at 30 days and 3 months***

At 30 days of age there was no difference in bodyweight, calorific intake, EE or glucose tolerance between offspring of obese and lean dams (data not shown).

At 3-months-of-age offspring did not differ between groups in bodyweight or body composition (percentage fat mass) as measured by bio-impedance (see Supplementary methods) or in calorific intake (data not shown).

Following ipGTT at 3 months, male OffOb showed an increase in the peak blood glucose concentration compared to OffCon although the area under the glucose curve was not different from OffCon (figure 3A, inset). In female OffOb, both peak blood glucose

## Maternal Obesity & Offspring Energy Expenditure

concentration after 15 minutes and AUC were elevated compared to OffCon. Maternal dietary supplementation of PDX resulted in lower peak blood glucose concentration in female OffObP at 15 minutes compared with OffOb (figure 3A).

There was no effect of maternal obesity on male or female 3-month-old offspring EE compared to controls. However, maternal dietary PDX supplementation in obese dams was associated with an increase in EE in male OffObP compared to OffOb during both day and night-time (Figure 3B). There was no effect of maternal PDX supplementation on EE in female OffOb (figure 3C).

Both male and female OffOb showed a significant reduction in respiratory exchange ratio compared to OffCon, which was not observed in female OffObP (figure 3B and C).

### ***Body Composition, Energy Balance and Glucose Tolerance at 6 months of age***

Bodyweight of 6-month-old male OffOb was increased compared to OffCon. Maternal dietary supplementation with PDX was associated with a reduction in body weight in male OffObP only ( $P<0.001$ , figure 4A).

The increase in male OffOb bodyweight was reflected in greater white adipose tissue mass (WAT) (Figure 4B) compared to OffCon, with an increase in the visceral fat pad mass (mesenteric fat mass [g]: OffOb:  $0.92 \pm 0.08$ ,  $n=6$  versus, OffCon  $0.65 \pm 0.03$ ,  $n=7$ ,  $P<0.05$ ). Maternal PDX supplementation prevented the rise in male offspring WAT mass and mesenteric fat mass secondary to maternal obesity (Figure 4B, mesenteric fat mass [g]: OffOb  $0.65 \pm 0.03$ ,  $n=7$ , versus OffObP  $0.51 \pm 0.06$ ,  $n=7$ ,  $P<0.05$ ).

Male OffOb had decreased brown adipose tissue (BAT) compared with OffCon (Figure 4C) when corrected for bodyweight. Maternal dietary PDX supplementation normalised BAT

weight relative to controls and resulted in male offspring with higher BAT weight compared to OffOb.

Both male and female OffOb demonstrated a greater peak glucose concentration in response to a glucose load (i.p.GTT, Figure 4D) and a greater AUC compared with OffCon. Maternal dietary PDX supplementation normalised offspring glucose profiles following the GTT (Figure 4D).

Maternal obesity resulted in lower EE in male OffOb during day and night compared with OffCon. Maternal PDX supplementation prevented the reduced EE associated with maternal obesity during both the active night-phase and the day-time rest-phase. There was no difference in EE between the female offspring at 6 months (Figure 4E and F).

### ***Obesogenic dietary challenge***

#### ***Body weight***

Male and female OffOb had greater body weight after three weeks' exposure to the obesogenic dietary challenge than similarly challenged OffCon. The exaggerated weight gain in both male and female OffOb on the obesogenic diet was prevented by maternal dietary PDX supplementation. (Figure 5A).

#### ***Energy Intake***

Calorific intake increased across all offspring groups following the obesogenic dietary challenge. Female, but not male, OffOb (figure 5B) increased calorific intake by 25% compared with OffCon fed the same hyper-calorific diet. Maternal dietary supplementation with PDX in the obese dams prevented the increased food intake in females on the high calorie diet.

### 309 *Energy expenditure*

310 Male and female offspring, in all experimental groups, showed decreased EE during their  
311 active (night-time) phase following the high fat dietary challenge. An observed reduction  
312 in EE after dietary challenge in adult male and female OffOb offspring compared to control  
313 was prevented by maternal PDX (Figure 5C) such that OffObP was similar to control.

314

### 315 *Respiratory Exchange Ratio*

316 The dietary challenge normalised respiratory exchange ratio across all groups, such that  
317 male and female OffOb no longer showed the reduction in RER observed at baseline  
318 (figure 5D).

319

### 320 ***Skeletal Muscle Mitochondrial DNA Copy number***

321 Mitochondrial DNA copy number was investigated as a potential determinant of the  
322 observed reduction in energy expenditure and glucose tolerance in OffOb. At 30 days-of-  
323 age, prior to any phenotypic change in the OffOb, MtDNA copy number ratio in male OffOb  
324 skeletal muscle was markedly increased compared to controls (Figure 6A). This was  
325 prevented by maternal PDX supplementation, such that OffObP males were similar to  
326 control. There was no significant effect of maternal diet on mitochondrial copy number  
327 ratio in female OffObP, although when sexes were combined there was a highly significant  
328 effect of maternal obesity on offspring skeletal muscle Mt/N ratio at 30 days, which was  
329 prevented by PDX.

330

### 331 ***Biomarkers of Brown Fat activation***

332 An increase in Dio2 mRNA expression in 6-month-old male OffOb, a gene encoding Type  
333 2 iodothyronine deiodinase involved in thermogenesis, was prevented by maternal  
334 dietary PDX supplementation. Pgc-1a mRNA expression (Peroxisome proliferator-  
335 activated receptor gamma coactivator 1-alpha) was increased in female OffOb and was

similarly prevented by maternal PDX (Figure 6B). Mitochondrial UCP-1, involved in non-shivering thermogenesis was up-regulated in male OffObP compared to OffCon (Figure 6B) but unaffected by maternal obesity alone.

### ***Gut Microbiota- faecal analysis of broad-spectrum gut bacterial species***

Maternal dietary PDX supplementation increased the percentage of *Bacteroides* in the male offspring bacterial population compared to OffCon at weaning (Figure 6C)

In 6-month-old offspring of obese dams the microbiota showed marked differences compared with controls (Figure 6C). Male and female OffOb demonstrated a higher percentage of *Eubacterium rectale-Clostridium coccoides* group compared with OffCon. There was no apparent influence of maternal PDX treatment on OffObP at 6 months.

## **DISCUSSION**

Here we report, in a mouse model, the influence of maternal obesity on offspring energy expenditure and the potential therapeutic benefit of maternal dietary intervention with the prebiotic polydextrose. Our main findings were firstly, that PDX improves glycaemic control and reproductive function in obese pregnancy, without affecting calorific intake or gestational weight gain; secondly, that maternal PDX treatment improves glucose homeostasis in both male and female offspring; and thirdly, that maternal PDX treatment prevents offspring weight gain, via sex specific changes in energy intake and energy expenditure. Lastly, maternal PDX supplementation provided protection against the effects of an obesogenic diet in adulthood.

***Maternal Phenotype***

Polydextrose has been shown to improve adult glucose metabolism (24) but not previously in pregnant women or obese pregnant animals. In the present study obese dams showed greatly improved glucose tolerance after supplementation with PDX. This was associated with an improvement in inflammatory cytokine profile in late gestation. Maternal glycaemia (and fetal hyperinsulinaemia) together with inflammatory mediators have been implicated in life-long obesity risk through the altered fetal hypothalamic neurodevelopment leading to disturbance of anabolic, adipogenic and neurotrophic pathways and permanent influences on metabolic and physiological development (25-27).

PDX also improved reproductive success in obese pregnant mice, with beneficial effects on fertility and litter size. Obesity perturbs the hypothalamic-pituitary-gonadal axis and ovarian cycle, reducing FSH and LH in the follicular and ovulatory phase whilst also shortening the luteal phase to reduce progesterone levels. It is possible, therefore, that PDX, either directly or indirectly, may influence reproductive hormones in gestation to improve reproductive capacity (28).

***Effect of maternal Obesity on Offspring Body Composition & Glucose tolerance***

Human cohort studies demonstrate that maternal overweight and obesity is associated with greater adiposity in offspring (5, 29). We found that male offspring were heavier with increased white adipose tissue mass and reduced BAT mass at 6 months of age. The impaired glucose tolerance observed in both males and females, at 3 months, antedates any observed changes in body composition (BIA) suggesting an alternative cause, potentially pancreatic beta cell dysfunction previously implicated in this model (30) or the early changes in mitochondrial function observed.

***Effect of maternal Obesity and PDX on Offspring Energy expenditure***

In this study we present novel evidence for the developmental programming of altered EE secondary to maternal diet-induced obesity, and prevention by maternal PDX supplementation. PDX influenced EE from as early as 3 months of age, preceding the subsequent changes in body composition, without affecting energy intake. Previously the scant literature in this area includes demonstration of reduced EE in 6-month-old infants born to overweight and obese mothers (31); in genetically altered mice following intrauterine exposure to gestational diabetes (32); and in 30 day old offspring of severely obesity rats (33). Changes in both RER and EE were associated with hepatic mitochondrial dysfunction, with reduced PGC-1 $\alpha$  mRNA expression, and impaired fatty acid oxidation (33). Taken together, these findings suggest impaired nutrient sensing and fuel switching in offspring of obese dams. Compromised fatty acid oxidation would be consistent with the development of a fatty liver phenotype which we have previously described in this rodent model (15, 17).

***Response to an obesogenic environment in adulthood: Energy intake on the obesogenic diet***

Female offspring of obese dams demonstrated hyperphagia secondary to maternal obesity only when exposed to obesogenic dietary challenge, suggesting programming of sex specific effects on food preference, and implicating mesolimbic reward pathways (34). Perinatal 'junk food' exposure similarly increases the preference for palatable diets in juvenile and adult rat offspring, and we previously reported reduced M $\mu$ -opioid receptor expression in the ventral tegmental area (VTA) of female 'junk-food' offspring only (35) (36). Moreover, we have previously reported in the offspring of obese rats, structural and functional deficits in neuronal development in the hypothalamic arcuate and paraventricular nucleus associated with leptin resistance and hyperphagia (49).



Prevention of female hyperphagia by maternal PDX supplementation, therefore, could imply protection of central neurotrophic development in the neonatal brain.

Male offspring of obese mice had lower energy expenditure than controls. In man, a blunted glucose-induced thermogenesis has been observed in obese individuals, increasing susceptibility to obesity when consuming diets rich in sugars (37-39). Since OffOb males were not obese at three months, a programmed deficit in diet-induced thermogenesis or central insulin resistance at the level of the hypothalamus could underlie the reduction in night time EE during the obesogenic dietary challenge. Reduced physical activity can also play a role in reduced EE, however, this is unlikely in the murine model employed here, since we have previously reported that male offspring of obese mice have a hyperactive ADHD-like phenotype (40)

### ***Mitochondrial biogenesis and activation***

The observed increase in mitochondrial DNA copy number in skeletal muscle at 30 days of age in OffOb males is consistent with early developmental exposure to maternal high glucose-induced ROS, secondary to maternal obesity, and could reflect compensatory mitochondrial biogenesis in response to a decline in mitochondrial function (21, 41, 42). Alternatively, the increase in MtDNA may be non-functional and a maladaptive response to oxidative stress (43) or hyperglycaemia (44), which can lead to an increase in tissue MtDNA and inflammation through activating of mTOR pathways and induction of TNF $\alpha$  (45). Either way, the data suggest an independent influence of maternal obesity on skeletal muscle mtDNA levels and hence mitochondrial function prior to the development of other metabolic defects, which might suggest a primary mechanism in the developmental programming due to maternal obesity.

443 ***Biomarkers of Brown Fat activation***

444 The increased expression Type 2 iodothyronine deiodinase (D2) which mediates adaptive  
445 thermogenesis in brown adipose tissue may reflect the increased sympathetic drive (46)  
446 previously described in this model (14, 47, 48). PDX may theoretically prevent this  
447 increased Dio2 gene expression by normalising hypothalamic development and  
448 sympathetic drive in OffOb (49). Indeed, others have shown that probiotics rescue  
449 neurogenesis and behavioural deficits in dysbiotic mice treated with antibiotics (50).

450

451 Mitochondrial UCP-1 expression was up-regulated in BAT of male offspring of obese dams  
452 treated with PDX compared to control offspring and may contribute to the increased  
453 energy expenditure observed. Prebiotics may increase thermogenic capacity in BAT by  
454 increasing UCP-1 expression (51) through altering microbiota and their by-products,  
455 short chain fatty acids, which can act as both energy source and receptor-mediated  
456 metabolic regulators of host energy metabolism involving processes such as hepatic  
457 gluconeogenesis and lipid metabolism via AMPK and PGC-1 $\alpha$  activation (52) .

458

459 Pgc1 $\alpha$  is the master regulator of mitochondrial biogenesis and linked to adaptive  
460 thermogenesis, following 'BAT activation'. Increased Pgc1 $\alpha$  expression in skeletal muscle  
461 of offspring of obese dams which was prevented by maternal PDX treatment appears  
462 counter-intuitive, as BAT activation would be expected to contribute to greater energy  
463 expenditure, if the observed increase in mtDNA were indeed functional. However, in  
464 addition to stimulating mitochondrial proliferation in skeletal muscle, PGC-1 $\alpha$  activation  
465 favours enhanced lipid- over carbohydrate-mediated mitochondrial respiration in  
466 skeletal muscle in mice, and leads to intrinsic mitochondrial adaptations in fatty acid-  
467 induced uncoupling and a reduction in mitochondrial superoxide production (53). This  
468 'fuel switching' is consistent with the observed reduction in RER, and thus increased lipid

oxidation, in offspring of obese mice and may represent a compensatory response to reduce ROS production, or a direct influence of the gut microbiota (52)

### ***Offspring microbiota profile***

Inheritable microbiota, passed from an obese mother to offspring during labour, may contribute to the modern patterns of human health and disease affecting gut barrier integrity and energy provision (54) but also maturation of the immune system (55), insulin sensitivity, energy expenditure and visceral adiposity (56). Indeed, we have previously implicated impaired innate immunity in offspring liver together with an increase in pro-inflammatory markers associated with NAFLD in offspring of obese mice (17). A recent landmark study demonstrated that transplanted gut microbiota from stool microbes of 2-week-old infants born to obese mothers increases inflammation and susceptibility to NAFLD in recipient germ-free mice (57).

### ***Prebiotic effects of polydextrose on offspring microbiota***

Maternal supplementation with PDX in obese pregnant mice resulted in increased abundance of *Bacteroides* compared to controls. Administration of prebiotics has previously been shown to improve pregnancy outcomes (58) and influence maternal transfer of microbiota and initial establishment of bifidobacteria in the infant (59). In obese humans an increase in bacteroides relative abundance is associated with weight-loss (60). We report a similar effect here, with maternal PDX intervention, in which obesity traits in the offspring were reduced associated with an increase in bacteroides relative abundance.

### **Conclusions**

In this study, evidence has been presented that diet-induced maternal obesity in the mouse results in reduced EE, glucose intolerance and increased bodyweight in 6 month

male offspring compared to controls. Moreover, following a 3-week obesogenic dietary challenge, offspring of obese dams had reduced energy expenditure, increased calorific intake and increased weight gain compared to controls. The offspring obesogenic phenotype is preceded by evidence of early mitochondrial damage and changes in the gut microbiota, which are prevented by maternal polydextrose. Polydextrose is a synthetic indigestible glucose polymer, classified as a dietary fibre and therefore, safe for use in pregnancy. However, there is currently a lack of high-quality scientific data on the use of polydextrose, or indeed other prebiotics, in pregnant or breastfeeding women. The present study supports the safety and efficacy of polydextrose supplementation in obese pregnancy.

*Supplementary information is available at International Journal of Obesity's website*

### **ACKNOWLEDGEMENTS**

*This work was supported by; a BBSRC industrial CASE studentship (BB/G017093/1) with Tate & Lyle PLC; Tommy's Charity; European Union 7th Framework Programme (FP7/2007-2013) project EarlyNutrition under grant agreement No. 289346, Medical Research Council UK; Newton Fund (Partnership Project grant) RCUK-CONACYT Research Partnerships MR/N029259/1. Polydextrose was provided by Tate & Lyle. This study also received support from the NIHR Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London. The views expressed in this publication are those of the author and not necessarily those of the NHS, the NIHR, or the Department of Health.*

### **CONFLICT OF INTEREST**

523 The authors declare no conflict of interest.

## 524 **References**

- 525 1. Poston L, Caleyachetty R, Cnattingius S, Corvalan C, Uauy R, Herring S, et al.  
526 Preconceptional and maternal obesity: epidemiology and health consequences. *Lancet*  
527 *Diabetes Endocrinol.* 2016;4(12):1025-36.
- 528 2. Godfrey KM, Reynolds RM, Prescott SL, Nyirenda M, Jaddoe VW, Eriksson JG, et al.  
529 Influence of maternal obesity on the long-term health of offspring. *Lancet Diabetes*  
530 *Endocrinol.* 2017;5(1):53-64.
- 531 3. Heslehurst N, Vieira R, Akhter Z, Bailey H, Slack E, Ngongalah L, et al. The  
532 association between maternal body mass index and child obesity: A systematic review  
533 and meta-analysis. *PLoS Med.* 2019;16(6):e1002817.
- 534 4. Scholtens DM, Kuang A, Lowe LP, Hamilton J, Lawrence JM, Lebenthal Y, et al.  
535 Hyperglycemia and Adverse Pregnancy Outcome Follow-up Study (HAPO FUS): Maternal  
536 Glycemia and Childhood Glucose Metabolism. *Diabetes Care.* 2019;42(3):381-92.
- 537 5. Dalrymple K, Thompson J, Begum S, *al. e.* Relationships of Maternal Body Mass  
538 Index and Plasma Biomarkers with Childhood Body Mass Index and Adiposity at 6 years;  
539 the Children of SCOPE study. *Int J Pead Ob.* 2019.
- 540 6. Hanson M, Barker M, Dodd JM, Kumanyika S, Norris S, Steegers E, et al.  
541 Interventions to prevent maternal obesity before conception, during pregnancy, and post  
542 partum. *Lancet Diabetes Endocrinol.* 2017;5(1):65-76.
- 543 7. do Carmo MM, Walker JC, Novello D, Caselato VM, Sgarbieri VC, Ouwehand AC, et  
544 al. Polydextrose: Physiological Function, and Effects on Health. *Nutrients.* 2016;8(9).
- 545 8. Ibarra A, Astbury NM, Olli K, Alhoniemi E, Tiihonen K. Effects of polydextrose on  
546 different levels of energy intake. A systematic review and meta-analysis. *Appetite.*  
547 2015;87:30-7.

- 548 9. Ibarra A, Astbury NM, Olli K, Alhoniemi E, Tiihonen K. Effect of Polydextrose on  
549 Subjective Feelings of Appetite during the Satiation and Satiety Periods: A Systematic  
550 Review and Meta-Analysis. *Nutrients*. 2016;8(1).
- 551 10. Ibarra A, Olli K, Pasman W, Hendriks H, Alhoniemi E, Raza GS, et al. Effects of  
552 polydextrose with breakfast or with a midmorning preload on food intake and other  
553 appetite-related parameters in healthy normal-weight and overweight females: An acute,  
554 randomized, double-blind, placebo-controlled, and crossover study. *Appetite*.  
555 2017;110:15-24.
- 556 11. Olli K, Salli K, Alhoniemi E, Saarinen M, Ibarra A, Vasankari T, et al. Postprandial  
557 effects of polydextrose on satiety hormone responses and subjective feelings of appetite  
558 in obese participants. *Nutr J*. 2015;14:2.
- 559 12. Airaksinen K, Jokkala J, Ahonen I, Auriola S, Kolehmainen M, Hanhineva K, et al.  
560 High-Fat Diet, Betaine, and Polydextrose Induce Changes in Adipose Tissue Inflammation  
561 and Metabolism in C57BL/6J Mice. *Mol Nutr Food Res*. 2018:e1800455.
- 562 13. Raza GS, Putaala H, Hibberd AA, Alhoniemi E, Tiihonen K, Makela KA, et al.  
563 Polydextrose changes the gut microbiome and attenuates fasting triglyceride and  
564 cholesterol levels in Western diet fed mice. *Sci Rep*. 2017;7(1):5294.
- 565 14. Samuelsson AM, Matthews PA, Argenton M, Christie MR, McConnell JM, Jansen EH,  
566 et al. Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity,  
567 hypertension, and insulin resistance: a novel murine model of developmental  
568 programming. *Hypertension*. 2008;51(2):383-92.
- 569 15. Oben JA, Mouralidarane A, Samuelsson AM, Matthews PJ, Morgan ML, McKee C, et  
570 al. Maternal obesity during pregnancy and lactation programs the development of  
571 offspring non-alcoholic fatty liver disease in mice. *J Hepatol*. 2010;52(6):913-20.
- 572 16. Oben JA, Patel T, Mouralidarane A, Samuelsson AM, Matthews P, Pombo J, et al.  
573 Maternal obesity programmes offspring development of non-alcoholic fatty pancreas  
574 disease. *Biochem Biophys Res Commun*. 2010;394(1):24-8.

## Maternal Obesity & Offspring Energy Expenditure

- 575 17. Muralidarane A, Soeda J, Visconti-Pugmire C, Samuelsson AM, Pombo J,  
576 Maragkoudaki X, et al. Maternal obesity programs offspring nonalcoholic fatty liver  
577 disease by innate immune dysfunction in mice. *Hepatology*. 2013;58(1):128-38.
- 578 18. Lagisz M, Blair H, Kenyon P, Uller T, Raubenheimer D, Nakagawa S. Little appetite  
579 for obesity: meta-analysis of the effects of maternal obesogenic diets on offspring food  
580 intake and body mass in rodents. *Int J Obes (Lond)*. 2015;39(12):1669-78.
- 581 19. Witaicenis A, Fruet AC, Salem L, Di Stasi LC. Dietary polydextrose prevents  
582 inflammatory bowel disease in trinitrobenzenesulfonic acid model of rat colitis. *J Med*  
583 *Food*. 2010;13(6):1391-6.
- 584 20. Stenman LK, Waget A, Garret C, Briand F, Burcelin R, Sulpice T, et al. Probiotic  
585 B420 and prebiotic polydextrose improve efficacy of antidiabetic drugs in mice. *Diabetol*  
586 *Metab Syndr*. 2015;7:75.
- 587 21. Malik AN, Czajka A. Is mitochondrial DNA content a potential biomarker of  
588 mitochondrial dysfunction? *Mitochondrion*. 2013;13(5):481-92.
- 589 22. Czajka A, Malik AN. Hyperglycemia induced damage to mitochondrial respiration  
590 in renal mesangial and tubular cells: Implications for diabetic nephropathy. *Redox Biol*.  
591 2016;10:100-7.
- 592 23. Rigottier-Gois L, Bourhis AG, Gramet G, Rochet V, Dore J. Fluorescent  
593 hybridisation combined with flow cytometry and hybridisation of total RNA to analyse  
594 the composition of microbial communities in human faeces using 16S rRNA probes. *FEMS*  
595 *Microbiol Ecol*. 2003;43(2):237-45.
- 596 24. Shimomura Y, Maeda K, Nagasaki M, Matsuo Y, Murakami T, Bajotto G, et al.  
597 Attenuated response of the serum triglyceride concentration to ingestion of a chocolate  
598 containing polydextrose and lactitol in place of sugar. *Biosci Biotechnol Biochem*.  
599 2005;69(10):1819-23.
- 600 25. Catalano PM, Shankar K. Obesity and pregnancy: mechanisms of short term and  
601 long term adverse consequences for mother and child. *BMJ*. 2017;356:j1.

## Maternal Obesity & Offspring Energy Expenditure

- 602 26. Roman AS, Rebarber A, Fox NS, Klauser CK, Istwan N, Rhea D, et al. The effect of  
603 maternal obesity on pregnancy outcomes in women with gestational diabetes. *J Matern*  
604 *Fetal Neonatal Med.* 2011;24(5):723-7.
- 605 27. Bo S, Menato G, Signorile A, Bardelli C, Lezo A, Gallo ML, et al. Obesity or diabetes:  
606 what is worse for the mother and for the baby? *Diabetes Metab.* 2003;29(2 Pt 1):175-8.
- 607 28. Al-Sobayil K, Zeitoun M, Abdel-Salam A. Effectiveness of a functional synbiotic  
608 syrup on pregnancy rate, neonatal birth weight and progesterone profile of oestrous-  
609 synchronized Najdi ewes *Journal of Food, Agriculture & Environment.* 2010;8(1):80-5.
- 610 29. Castillo-Laura H, Santos IS, Quadros LC, Matijasevich A. Maternal obesity and  
611 offspring body composition by indirect methods: a systematic review and meta-analysis.  
612 *Cad Saude Publica.* 2015;31(10):2073-92.
- 613 30. Taylor PD, McConnell J, Khan IY, Holemans K, Lawrence KM, Asare-Anane H, et al.  
614 Impaired glucose homeostasis and mitochondrial abnormalities in offspring of rats fed a  
615 fat-rich diet in pregnancy. *Am J Physiol Regul Integr Comp Physiol.* 2005;288(1):R134-9.
- 616 31. Rising R, Lifshitz F. Lower energy expenditures in infants from obese biological  
617 mothers. *Nutr J.* 2008;7:15.
- 618 32. Lau SM, Lin S, Stokes RA, Cheng K, Baldock PA, Enriquez RF, et al. Synergistic  
619 effects of genetic beta cell dysfunction and maternal glucose intolerance on offspring  
620 metabolic phenotype in mice. *Diabetologia.* 2011;54(4):910-21.
- 621 33. Borengasser SJ, Lau F, Kang P, Blackburn ML, Ronis MJ, Badger TM, et al. Maternal  
622 obesity during gestation impairs fatty acid oxidation and mitochondrial SIRT3 expression  
623 in rat offspring at weaning. *PLoS One.* 2011;6(8):e24068.
- 624 34. Ong ZY, Muhlhausler BS. Maternal "junk-food" feeding of rat dams alters food  
625 choices and development of the mesolimbic reward pathway in the offspring. *FASEB J.*  
626 2011;25(7):2167-79.



## Maternal Obesity & Offspring Energy Expenditure

- 627 35. Gugusheff JR, Bae SE, Rao A, Clarke IJ, Poston L, Taylor PD, et al. Sex and age-  
628 dependent effects of a maternal junk food diet on the mu-opioid receptor in rat offspring.  
629 Behav Brain Res. 2016;301:124-31.
- 630 36. Bayol SA, Farrington SJ, Stickland NC. A maternal 'junk food' diet in pregnancy and  
631 lactation promotes an exacerbated taste for 'junk food' and a greater propensity for  
632 obesity in rat offspring. Br J Nutr. 2007;98(4):843-51.
- 633 37. Bogardus C, Lillioja S, Mott D, Zawadzki J, Young A, Abbott W. Evidence for  
634 reduced thermic effect of insulin and glucose infusions in Pima Indians. J Clin Invest.  
635 1985;75(4):1264-9.
- 636 38. Schutz Y, Bessard T, Jequier E. Diet-induced thermogenesis measured over a  
637 whole day in obese and nonobese women. Am J Clin Nutr. 1984;40(3):542-52.
- 638 39. Golay A, Schutz Y, Felber JP, Jallut D, Jequier E. Blunted glucose-induced  
639 thermogenesis in 'overweight' patients: a factor contributing to relapse of obesity. Int J  
640 Obes. 1989;13(6):767-75.
- 641 40. Fernandes C, Grayton H, Poston L, Samuelsson AM, Taylor PD, Collier DA, et al.  
642 Prenatal exposure to maternal obesity leads to hyperactivity in offspring. Mol Psychiatry.  
643 2012;17(12):1159-60.
- 644 41. Rooney JP, Ryde IT, Sanders LH, Howlett EH, Colton MD, Germ KE, et al. PCR based  
645 determination of mitochondrial DNA copy number in multiple species. Methods Mol Biol.  
646 2015;1241:23-38.
- 647 42. Al-Kafaji G, Sabry MA, Bakhiet M. Increased expression of mitochondrial DNA-  
648 encoded genes in human renal mesangial cells in response to high glucose-induced  
649 reactive oxygen species. Mol Med Rep. 2016;13(2):1774-80.
- 650 43. Masser DR, Clark NW, Van Remmen H, Freeman WM. Loss of the antioxidant  
651 enzyme CuZnSOD (Sod1) mimics an age-related increase in absolute mitochondrial DNA  
652 copy number in the skeletal muscle. Age (Dordr). 2016;38(4):323-33.

- 653 44. Malik AN, Parsade CK, Ajaz S, Crosby-Nwaobi R, Gnudi L, Czajka A, et al. Altered  
654 circulating mitochondrial DNA and increased inflammation in patients with diabetic  
655 retinopathy. *Diabetes Res Clin Pract.* 2015;110(3):257-65.
- 656 45. Picca A, Lezza AMS, Leeuwenburgh C, Pesce V, Calvani R, Bossola M, et al.  
657 Circulating Mitochondrial DNA at the Crossroads of Mitochondrial Dysfunction and  
658 Inflammation During Aging and Muscle Wasting Disorders. *Rejuvenation Res.*  
659 2018;21(4):350-9.
- 660 46. de Jesus LA, Carvalho SD, Ribeiro MO, Schneider M, Kim SW, Harney JW, et al. The  
661 type 2 iodothyronine deiodinase is essential for adaptive thermogenesis in brown adipose  
662 tissue. *J Clin Invest.* 2001;108(9):1379-85.
- 663 47. Samuelsson AM, Clark J, Rudyk O, Shattock MJ, Bae SE, South T, et al. Experimental  
664 hyperleptinemia in neonatal rats leads to selective leptin responsiveness, hypertension,  
665 and altered myocardial function. *Hypertension.* 2013;62(3):627-33.
- 666 48. Samuelsson AM, Morris A, Igosheva N, Kirk SL, Pombo JM, Coen CW, et al. Evidence  
667 for sympathetic origins of hypertension in juvenile offspring of obese rats. *Hypertension.*  
668 2010;55(1):76-82.
- 669 49. Kirk SL, Samuelsson AM, Argenton M, Dhonye H, Kalamatianos T, Poston L, et al.  
670 Maternal obesity induced by diet in rats permanently influences central processes  
671 regulating food intake in offspring. *PLoS One.* 2009;4(6):e5870.
- 672 50. Mohle L, Mattei D, Heimesaat MM, Bereswill S, Fischer A, Alutis M, et al. Ly6C(hi)  
673 Monocytes Provide a Link between Antibiotic-Induced Changes in Gut Microbiota and  
674 Adult Hippocampal Neurogenesis. *Cell Rep.* 2016;15(9):1945-56.
- 675 51. Rodriguez VM, Portillo MP, Pico C, Macarulla MT, Palou A. Olive oil feeding up-  
676 regulates uncoupling protein genes in rat brown adipose tissue and skeletal muscle. *Am J*  
677 *Clin Nutr.* 2002;75(2):213-20.

- 678 52. den Besten G, van Eunen K, Groen AK, Venema K, Reijngoud DJ, Bakker BM. The  
679 role of short-chain fatty acids in the interplay between diet, gut microbiota, and host  
680 energy metabolism. *J Lipid Res.* 2013;54(9):2325-40.
- 681 53. Hoeks J, Arany Z, Phielix E, Moonen-Kornips E, Hesselink MK, Schrauwen P.  
682 Enhanced lipid-but not carbohydrate-supported mitochondrial respiration in skeletal  
683 muscle of PGC-1alpha overexpressing mice. *J Cell Physiol.* 2012;227(3):1026-33.
- 684 54. Zhu MJ, Du M, Ford SP. Impacts of maternal obesity on placental and gut  
685 inflammation and health. *J Anim Sci.* 2013.
- 686 55. Myles IA, Fontecilla NM, Janelins BM, Vithayathil PJ, Segre JA, Datta SK. Parental  
687 dietary fat intake alters offspring microbiome and immunity. *J Immunol.*  
688 2013;191(6):3200-9.
- 689 56. Stolarczyk E, Vong CT, Perucha E, Jackson I, Cawthorne MA, Wargent ET, et al.  
690 Improved insulin sensitivity despite increased visceral adiposity in mice deficient for the  
691 immune cell transcription factor T-bet. *Cell Metab.* 2013;17(4):520-33.
- 692 57. Soderborg TK, Clark SE, Mulligan CE, Janssen RC, Babcock L, Ir D, et al. The gut  
693 microbiota in infants of obese mothers increases inflammation and susceptibility to  
694 NAFLD. *Nat Commun.* 2018;9(1):4462.
- 695 58. Laitinen K, Poussa T, Isolauri E, Nutrition AMI, Intestinal Microbiota G. Probiotics  
696 and dietary counselling contribute to glucose regulation during and after pregnancy: a  
697 randomised controlled trial. *Br J Nutr.* 2009;101(11):1679-87.
- 698 59. Collado MC, Cernada M, Bauerl C, Vento M, Perez-Martinez G. Microbial ecology  
699 and host-microbiota interactions during early life stages. *Gut Microbes.* 2012;3(4):352-  
700 65.
- 701 60. Ley RE, Turnbaugh PJ, Klein S, Gordon JI. Microbial ecology: human gut microbes  
702 associated with obesity. *Nature.* 2006;444(7122):1022-3.
- 703

**FIGURE LEGENDS**

**Figure 1. Schematic representation of the experimental design.** Female dams were fed either an obesogenic (n=34) or a control diet (n=18). Following successful mating a subgroup of obese dams were supplemented with 5 % PDX (n=12) in the drinking water. All offspring were weaned on to control diet and were followed up to 6 months. 1 male and 1 female was studied at each timepoint: 30 days, 3 months and 6 months. At three months after recording baseline characteristics, 1 male and 1 female from each litter (n=10) were exposed to the obesogenic diet for three weeks and reassessed.

**Figure 2. Maternal characteristics (A)** Gestational bodyweights and **(B)** calorific intake during gestation in obese (Ob) and obese supplemented with PDX (ObP) dams (n=6-7). **(C)** Response to a glucose tolerance test (GTT) and the respective area under the curve (AUC) on GD16 **(D)** Litter size and **(E)** Birthweight in control (Con) obese (Ob) and obese supplemented with PDX (ObP) dams (n=8-16) **(F)** Cytokine profile from late gestation obtained by 2 samples of pooled serum samples (n=5 per pool) from dams at day 16. Data are expressed as mean  $\pm$  SEM. \*represents  $P<0.05$ , \*\* represents  $P<0.01$ , comparison with the obese group.

**Figure 3. Offspring Phenotype at 3 months (A)** Response to a glucose tolerance test (GTT) and the respective area under the curve (AUC, inset). Average energy expenditure (EE) and RER during daytime or night-time (as indicated) in male **(B)** and female **(C)** offspring of control (OffCon), obese (OffOb) and obese dams supplemented with PDX (OffObP) at 3 months of age, n=6-15; Data are expressed as mean  $\pm$  SEM. \* represents  $P<0.05$ ; \*\* represents  $P<0.01$  versus OffCon. Hash symbol # represents  $P<0.05$  male OffObP vs OffOb.

**Figure 4. Offspring Phenotype at 6 months** Average **(A)** bodyweight **(B)** weight of white adipose tissue (WAT) and **(C)** brown adipose tissue weight corrected for bodyweight **(D)** GTT with AUC inset **(E)** Day-time Energy expenditure and **(F)** night-time energy expenditure in male and female offspring of control (OffCon), obese (OffOb) and obese dams supplemented with PDX (OffObP) at 6 months of age, (n=6-7), Data are expressed as mean  $\pm$  SEM. \* represents  $P<0.05$ , \*\* represents  $P<0.01$ , \*\*\* represents  $P<0.001$ .

**Figure 5. Obesogenic Dietary Challenge.** Offspring Phenotype at 3 months after 3 weeks on the obesogenic diet **(A)** Bodyweights and **(B)** average daily calorific intake **(C)** energy expenditure and **(D)** RER in male and female offspring of control (OffCon), obese (OffOb) and obese dams supplemented with PDX (OffObP) at 3 months of age and after three-weeks exposure to obesogenic diet, n=7-10. Exposure to obesogenic diet (OD) and maternal diet significantly accounted for variation (two-way ANOVA); Data are expressed as mean  $\pm$  SEM. \* represents  $P<0.05$ , \*\* represents  $P<0.01$ , \*\*\* represents  $P<0.001$ .

**Figure 6. Potential mechanisms** **(A)** MtDNA copy number in skeletal muscle from male and female offspring of control (OffCon), obese (OffOb) and obese dams supplemented with PDX (OffObP) at 30 days of age, (n= 5-6) **(B)** mRNA expression of brown adipose tissue biomarkers of BAT activity at 30 days of age. **(C)** Percentage of *bacteroides* (Bac+) and **(D)** *Eubacterium rectale-Clostridium coccoides* (Erec+) in bacteria cells (EUB+) identified in faecal samples from male and female offspring of control (OffCon), obese (OffOb) and obese dams supplemented with PDX (OffObP) at **(C)** weaning and **(D)** 6 months of age (n=6). **(E)** Representative FACS plot. EUB+ and gated cells. FL1 histogram, green fluorescence is the total number of bacteria hybridising with the EUB 338-FITC probe. FL4 histogram, red fluorescence, shows the proportion of cells targeted by the

group Cy5-probe in the sample. Data are expressed as mean  $\pm$  SEM. \* represents  $P < 0.05$ ;  
 \*\* represents  $P < 0.01$ , \*\*\* represents  $P < 0.001$ .

### Supplementary Methods

#### *Animal husbandry and diets*

Female C57BL/6J mice were maintained under controlled conditions (22°C, 12-hour light/dark cycle) and fed either a standard chow diet (RM1, Special Dietary Services, UK, 7% simple sugars, 3% fat, 50% polysaccharide, 15% protein [w/w] energy 3.5 kcal/g; n=18) or a semisynthetic obesogenic diet (#824053, Special Dietary Service, 10% simple sugars, 20% animal lard 28% polysaccharide, 23% protein [w/w], Special Dietary Services, energy 4.5 kcal/g, n=46). The obesogenic diet was supplemented by *ad libitum* access to sweetened condensed milk, fortified to control levels with micronutrient mineral mix (AIN93G, Special Dietary Services, UK). The combined macronutrient and calorific composition of the highly palatable obesogenic diet (based on intake) was approximately 16% fat, 33% simple sugars, 15% protein, total energy 4.0 kcal/g. Diets were introduced 6 weeks prior to mating and throughout gestation and lactation, as previously described by Samuelsson *et al* (2008) (14). All females included in the study were proven breeders with one previous successful pregnancy. Following successful mating, with proven breeders, indicated by the presence of a copulation plug, a sub-group of obesogenic diet-fed dams were randomly assigned to the same obesogenic diet supplemented with PDX (5% w/v) in the drinking water throughout gestation and lactation (n= 34) leading to three groups; control (Con); obese (Ob) and obese +5% (w/v) PDX, ObP, *Figure 1*). This concentration of PDX has previously proven efficacious in reducing insulin resistance in adult rats without toxicity and without altering calorific intake (19).

All pups were weighed 48 hours after delivery (to avoid pup rejection), and each litter was then standardised to 3 males and 3 females where possible. Litters with less than four pups were not included in the study.

### ***Indirect Calorimetry***

At each time point, 6 age-matched animals were weighed, and then placed in the open circuit indirect calorimetry cages (CaloCages) for 24 hours of acclimatisation, followed by 24 hours of experimental recording. Data were recorded automatically every 20 minutes and hourly means calculated.

### ***Mitochondrial DNA Copy Number in Offspring Skeletal Muscle***

2ul sample DNA diluted to 10ng/ul was loaded in triplicate alongside a 5-point standard curve consisting of primer-specific amplicons of known copy number.

### ***Detailed Glucose tolerance test***

In order to determine whole body glucose tolerance, an intra-peritoneal glucose tolerance test (ipGTT) was performed using AlphaTRAK® Glucose meter (Abbott Animal health), which has been specially designed for and validated in mice and rats. Following an overnight fast, a cream containing lidocaine (2.5%) and prilocaine (2.5%) (EMLA cream 5%, AstraZeneca, UK) was applied as a topical anaesthetic to the tail. Once the analgesia had taken effect, fasting tail venous blood glucose was measured. Animals were then injected via the intraperitoneal (i.p.) route of administration with a glucose load (1 gram/kg) of glucose solution (10% glucose). Measurements of the blood glucose were taken at 15, 30, 60 and 120 minutes after the glucose injection using the glucose meter. Measurements were taken at each point while the animals were conscious and semi-restrained. Glucose tolerance tests were performed on the dams at gestational day 16 (GD16) and on the offspring at 30, 90 and 180 days of age.

812 ***Biochemical analysis (Cytokine Profile)***

813 Briefly, the Proteome Profiler Mouse Adipokine Array allows the simultaneous  
814 measurement of relative expression levels of 38 mouse adipokines. Capture and control  
815 antibodies were spotted in duplicate on nitrocellulose membranes. Serum samples were  
816 diluted, mixed with a biotinylated detection antibody, and incubated overnight with the  
817 Proteome Profiler Mouse Adipokine Array. The next morning the membrane was washed  
818 in order to remove unbound material. Streptavidin-HRP and chemiluminescent detection  
819 reagents were applied allowing the production of a signal at each capture spot  
820 corresponding to the amount of protein bound. The density of each protein was measured  
821 with Image J.

822

823 **Analysis with flow cytometry**

824 The analysis was performed with flow cytometry as described by Rigottier-Gois et al (23).  
825 Cells were pelleted and resuspended in PBS for data acquisition by flow cytometry (HTS  
826 Fortessa, Becton Dickinson, USA).

827

828 A total of 20 000 events EUB 338-FITC positives were stored in list mode files. Subsequent  
829 analyses were conducted using FlowJo software (Tree Star, USA). Cell enumeration was  
830 performed by combining, in one hybridisation tube, one group Cy5-probe with the EUB  
831 338-FITC probe. An FL1 histogram (green fluorescence) was used to evaluate the total  
832 number of bacteria hybridising with the EUB 338-FITC probe. A gate was designed in this  
833 histogram representing the total number of bacterial cells in the sample and was used to  
834 build an FL4 histogram (red fluorescence) to directly estimate the proportion of cells  
835 targeted by the group Cy5-probe in the sample. The proportion of cells was corrected by  
836 eliminating background fluorescence, which was measured using the negative control  
837 NON 338-Cy5 probe. Results were expressed as cells hybridising with the group-Cy5



## Maternal Obesity & Offspring Energy Expenditure

838 probe as a proportion of the total bacteria hybridising with the EUB 338-FITC bacteria  
839 domain probe or normalised to the basal conditions as indicated.  
840

## Supplementary tables

Diet	Control (RM1)	Obesogenic Diet (Pellets)	Obesogenic Diet (Sweet Condensed milk)
Protein (%)	15	23	7.8
Total Carbohydrate (%)	61.73	38.83	55.3
Polysaccharides (%)	57.68	28.34	-
Simple Sugars (%)	7	10.49	55.3
Fat (%)	2.78	22.21	8.1
Soya oil (%)	-	4.32	-
Lard (%)	-	17.89	5.6
Corn oil (%)	2.78		
Crude Fibre (%)	4.65	6.17	traces
Energy (kcal/g)	3.5	4.5	3.3

**Table S 1.** Composition of the experimental diets presented as percentages by weight. The obesogenic diet consisted of both pellets and sweetened condensed milk. Due to varying moisture and nitrogen free extract content, rows will not sum to 100%.

Primers	Accession number	Sequences
mouse mitochondrion complete genome	NC_005089.1	<b>Forward primer:</b> 5'-CTAGAAACCCCGAAACCAAA - 3' <b>Reverse primer:</b> 5'-CCAGCTATCACCAAGCTCGT-3'
mouse $\beta$ 2M (beta-2 microglobulin)	NC_000068.8	<b>Forward primer:</b> 5'-CTAGAAACCCCGAAACCAAA - 3' <b>Reverse primer:</b> 5'-CCAGCTATCACCAAGCTCGT-3'

**Table S2** Oligonucleotide primer sequences used to determine mitochondrial copy number in skeletal muscle (tibialis anterior) samples.

Gene	Forward Primer	Reverse Primer
<i>Cyclophilin B</i>	TGGAGAGCACCAAGACAGACA	TGCCGGAGTCGACAATGAT
<i>Dio2</i>	CCTACAAACAGGTTAAACTGGG	CTCTGCACTGGCAAAGTC
<i>Pgc1a</i>	TGAAAGGGCCAAACAGAGAGA	TAAATCACACGGCGCTCTT
<i>Ucp1</i>	AATACTGGCAGATGACGTCC	TTACCACATCCACTGGAGAG
<i>Zic1</i>	CACATGAAGGTCCATGAGTCC	GGGTTGTCTGTTGTGGGAG

**Table S3** Semi-quantitative real-time PCR primers and sequences for BAT tissue samples.

Probe	Sequence	Target	Label 5'
EUB 338 pB-00159	GCTGCCTCCCGTAGGAGT	Domain Bacteria	FITC
NON 338 pB-00243	ACATCCTACGGGAGGC	Negative probe	Cy5
Bac 303 pB-00031	CCAATGTGGGGGACCTT	<i>Bacteroides</i>	Cy5
Erec 482 pB-00963	GCTTCTTAGTCARGTACCG	<i>Clostridium coccoides- Eubacterium rectale</i>	Cy5
Lab 158 pB-03928	GGTATTAGCAYCTGTTTCCA	<i>Lactobacillus-Streptococcus group</i>	Cy5
Bif 164 pB-00037	CATCCGGCATTACCACCC	<i>Bifidobacterium</i>	Cy5

**Table S4.** 16S rRNA-targeted oligonucleotide probes, and targets for rRNA dot-blot hybridisation (Panel of group- and species-specific 16S rRNA-targeted oligonucleotide probes)